

Note on Integrability of Marginally Deformed ABJ(M) Theories

Song He,^{a,b} Jun-Bao Wu^c

^a*State Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Science, No.55 Zhongguancun East Road, Beijing 100190, P. R. China*

^b*Kavli Institute for Theoretical Physics China, CAS, No.55 Zhongguancun East Road, Beijing 100190, P. R. China*

^c*Institute of High Energy Physics, and Theoretical Physics Center for Science Facilities, Chinese Academy of Sciences, 19B YuquanLu, Beijing 100049, P.R. China*

E-mail: hesong@itp.ac.cn, wujb@ihep.ac.cn

ABSTRACT: We study the anomalous dimensions of operators in the scalar sector of β -deformed ABJ(M) theories. We show that the anomalous dimension matrix at two-loop order gives an integrable Hamiltonian acting on an alternating $SU(4)$ spin chain with the spins at odd lattice sides in the fundamental representation and the spins at even lattices in the anti-fundamental representation. We get a set of β -deformed Bethe ansatz equations which give the eigenvalues of Hamiltonian of this deformed spin chain system. Based on our computations, we also extend our study to non-supersymmetric three-parameter γ -deformation of ABJ(M) theories and find that the corresponding Hamiltonian is the same as the one in β -deformed case at two-loop level in the scalar sector.

KEYWORDS: Gauge-gravity correspondence, Bethe ansatz, Supersymmetric gauge theory

ARXIV EPRINT: [1302.2208](https://arxiv.org/abs/1302.2208)

Contents

1	Introduction	1
2	β-deformation of superconformal Chern-Simons-matter theory	3
3	Two-loop anomalous dimensions in the scalar sector	6
4	The R matrices of deformed spin chain	9
5	Eigenvalues of deformed spin chain Hamiltonian and Bethe ansatz equations	12
6	The non-supersymmetric three-parameter γ-deformation of ABJ(M) theory	13
7	Conclusion and Discussions	14
A	Hamiltonian from deformed R-matrices	15

1 Introduction

There were great achievements on integrable structure in both sides of AdS/CFT correspondence in the last decade [1]. The best studied case is the integrability in the correspondence between four-dimensional $\mathcal{N} = 4$ super-Yang-Mills theory and the type IIB superstring theory on $AdS_5 \times S^5$ [2–4]. In the field theory side, it was found that, first at lower loop level, the anomalous dimension matrix coincides with Hamiltonian of certain integrable spin chain [5–7]. The all-loop Bethe ansatz equations [8] were later proposed and they can be obtained from a S-matrix for the spin chain [9], though we do not know the corresponding Hamiltonian of the spin chain at all-loop level. In the string theory side, infinity number of conserved charges on the worldsheet of Green-Schwarz superstring moving in $AdS_5 \times S^5$ were constructed [10]. The integrable structure is a very powerful tool which enables us to compute, as an example, the cusp anomalous dimension for arbitrary value of the 't Hooft coupling in the planar limit [11, 12].

Such integrable structure was also found for the recently proposed duality [13, 14] between three-dimensional $\mathcal{N} = 6$ Chern-Simons-matter theory and type IIA theory on $AdS_4 \times CP^3$ [15]–[42] (for reviews see [43, 44]). The dynamics in this case is more complicated and richer than the previous one, partly due to the fact that we now have less supersymmetries. As an example, there is still a to-be-determined function in the dispersion relation of the magnon [24, 25, 45]. In the $\mathcal{N} = 4$ super Yang-Mills case, this function

is trivial due to the fact that the theory is self-dual under the S-duality transformation [46].

With the success of integrability in mind, it should be with great value to generalize the above studies to case with less supersymmetries. The theory in which the anomalous dimensions of gauge invariant operators are related to an integrable Hamiltonian seems to be quite rare. If we perform generic marginal deformations of the $\mathcal{N} = 4$ super Yang-Mills theory [47] or ABJM theory, the obtained theory seems usually not to keep these integrable structure in the above sense, even when some supersymmetries and conformal symmetry are preserved. The β - and γ -deformations of $\mathcal{N} = 4$ super Yang-Mills theory are quite special since they preserve this remarkable integrable structure [48]-[51] and their gravity dual can be obtained through a certain solution generating technique [52]. Further studies of this integrability can be found in [53]-[73]. These marginal deformations are special also because they can be expressed elegantly using a star product which produces a certain phase factor for each interaction term in the Lagrangian. The solution generating transformation in the gravity side can be constructed by T-duality-shift-T-duality transformations in string theory [52]. The understand of the gauge-gravity correspondence in this case was improved in [74]. The β - and γ - deformed ABJM theories and their gravity duals were also studied in [74]. Some classical string solutions in these deformed backgrounds of type IIA string theory have been studied in [75]-[78]. The aim of this paper is to explore the integrable structure in the field theory side of these deformed AdS_4/CFT_3 correspondence.

We begin with the scalar sector of the β -deformed ABJ(M) theory which has $\mathcal{N} = 2$ supersymmetries. We compute the anomalous dimensions of these operators at the two-loop level in the planar limit and for all operators with length larger than 2. We express the result as a Hamiltonian of an alternating $SU(4)$ spin chain. Comparing with the undeformed case [15–17], we find that, in the Hamiltonian, only the terms from the interaction terms with six scalars are deformed. Though the interaction terms with two scalars and two fermions are also deformed, their contributions to the Hamiltonian coincide with the undeformed case. We also find that the Hamiltonian in non-supersymmetric three-parameter γ -deformed ABJ(M) theory is the same as the one of the β -deformed theory, at two-loop level in the scalar sector. We expect the differences will appear in other sectors and/or at higher loop order. As in [50], we deform the R-matrices constructed in [15, 16] by introducing suitable phase factors. We show that the obtained transfer matrices will produce essentially the Hamiltonian from the perturbative computations in the Chern-Simons-matter theories. This result shows that the Hamiltonian is integrable. By diagonalizing the transfer matrices, we obtain the Bethe ansatz equations and the eigenvalues of the Hamiltonian.

The organization of the remaining parts of this paper is as follows. In section 2 we briefly review β -deformation of ABJM theory. In section 3, we compute the two-loop corrections to the anomalous dimensions of operators in the scalar sector in both ABJM and ABJ theories. In section 4, we constructed the R matrices after deformation and show that the Hamiltonian obtained in section 2 is integrable. Based on these results, we derive the eigenvalues of the Hamiltonian of the deformed spin chain system in section 5. A brief discussion on non-supersymmetric three-parameter γ -deformation is put in

section 6. Section 7 is devoted to conclusions and discussions. We put some details of the computations in section 4 in the Appendix.

2 β -deformation of superconformal Chern-Simons-matter theory

The ABJM theory [13] is three dimensional $\mathcal{N} = 6$ supersymmetric Chern-Simons-matter theory. The gauge group of this theory is $U(N) \times U(N)$, and the Chern-Simons levels of these two subgroups are k and $-k$, respectively. The matter fields are four complex scalars $Y^I, I = 1, 2, 3, 4$ and four fermions $\Psi_I, I = 1, 2, 3, 4$ in the (N, \bar{N}) representation. The action of this theory is *

$$S = \int d^3x (L_{CS} + L_{kin.} - V_F - V_B), \quad (2.1)$$

with

$$L_{CS} = \frac{k}{4\pi} \epsilon^{\mu\nu\rho} \text{Tr}(A_\mu \partial_\nu A_\rho + \frac{2i}{3} A_\mu A_\nu A_\rho) - \frac{k}{4\pi} \epsilon^{\mu\nu\rho} \overline{\text{Tr}}(\bar{A}_\mu \partial_\nu \bar{A}_\rho + \frac{2i}{3} \bar{A}_\mu \bar{A}_\nu \bar{A}_\rho), \quad (2.2)$$

$$L_{kin.} = \frac{1}{2} \overline{\text{Tr}}(-(D_\mu Y)^\dagger D^\mu Y^I + i \Psi^{\dagger I} \gamma_\mu D^\mu \Psi_I) + \frac{1}{2} \text{Tr}(-D_\mu Y^I (D^\mu Y)_I^\dagger + i \Psi_I \gamma_\mu D^\mu \Psi^{\dagger I}), \quad (2.3)$$

$$V_B = -\frac{1}{3} \left(\frac{2\pi}{k} \right)^2 \overline{\text{Tr}} \left[Y_I^\dagger Y^J Y_J^\dagger Y^K Y_K^\dagger Y^I + Y_I^\dagger Y^I Y_J^\dagger Y^J Y_K^\dagger Y^K + 4Y_I^\dagger Y^J Y_K^\dagger Y^I Y_J^\dagger Y^K - 6Y_I^\dagger Y^I Y_J^\dagger Y^K Y_K^\dagger Y^J \right], \quad (2.4)$$

$$V_F = \frac{2\pi i}{k} \overline{\text{Tr}} \left[Y_I^\dagger Y^I \Psi^{\dagger J} \Psi_J - 2Y_I^\dagger Y^J \Psi^{\dagger I} \Psi_J + \epsilon^{IJKL} Y_I^\dagger \Psi_J Y_K^\dagger \Psi_L \right] - \frac{2\pi i}{k} \text{Tr} \left[Y^I Y_I^\dagger \Psi_J \Psi^{\dagger J} - 2Y^I Y_J^\dagger \Psi_I \Psi^{\dagger J} + \epsilon_{IJKL} Y^I \Psi^{\dagger J} Y^K \Psi^{\dagger L} \right]. \quad (2.5)$$

The covariant derivatives are:

$$D_\mu Y^I = \partial_\mu Y^I + i A_\mu Y^I - i Y^I \bar{A}_\mu, D_\mu Y_I^\dagger = \partial_\mu Y_I^\dagger + i \bar{A}_\mu Y_I^\dagger - i Y_I^\dagger \bar{A}_\mu, \quad (2.6)$$

$$D_\mu \Psi_I = \partial_\mu \Psi_I + i A_\mu \Psi_I - i \Psi_I \bar{A}_\mu, D_\mu \Psi^{\dagger I} = \partial_\mu \Psi^{\dagger I} + i \bar{A}_\mu \Psi^{\dagger I} - i \Psi^{\dagger I} \bar{A}_\mu. \quad (2.7)$$

In the following part, we will discuss the β deformation of the theory following the convention given by [74]. The deformed theory will preserve three dimensional $\mathcal{N} = 2$ supersymmetries. The β -deformation can be performed by replacing all of the ordinary product fg of two fields f and g in the Lagrangian by the following star product:

$$f * g = e^{i\pi\gamma(Q_1^f Q_2^g - Q_2^f Q_1^g)} fg, \quad (2.8)$$

*We follow the convention of [16] closely.

Table 1. $U(1)^2$ charges of the scalars of the ABJM theory used for β -deformation.

	Y^1	Y^2	Y^3	Y^4
$U(1)_1$	$+\frac{1}{2}$	$-\frac{1}{2}$	0	0
$U(1)_2$	0	0	$\frac{1}{2}$	$-\frac{1}{2}$

where $Q_i^f, i = 1, 2$ are two global $U(1)$ charges carrying by the field f and γ is a real deformation parameter. [†] For the β -deformation of ABJM theory, we choose the charges for the scalars as in table 1. The fermionic super-partner $\psi^{\dagger I}$ carries the same charge as Y^I , and the gauge field is neutral under these symmetries.

It is easy to see that this star product is associative and for several fields F_1, \dots, F_n , we have

$$F_1 * \dots * F_n = e^{i\pi\gamma \sum_{i < j} (Q_1^{F_i} Q_2^{F_j} - Q_2^{F_i} Q_1^{F_j})} F_1 \dots F_n. \quad (2.9)$$

This deformation just adds phase factors according to the above equation to the interaction terms in the Lagrangian. One can see from the above rule for the star product that the deformation does not change the kinetic terms of the Lagrangian. In the superfield formulism, the deformation will only deform the superpotential W in the following manner given in [74]:

$$W \rightarrow W^{\text{deformed}} = \frac{4\pi}{k} \text{Tr} \left(e^{-i\pi\gamma/2} Y^1 Y_3^\dagger Y^2 Y_4^\dagger - e^{i\pi\gamma/2} Y^1 Y_4^\dagger Y^2 Y_3^\dagger \right), \quad (2.10)$$

where one can go back to the original quartic superpotential by setting deformation parameter γ to be zero.

For later use, we now give the interaction terms after the deformation. One can see that only V_B and V_F will be deformed. By a bit calculations, we found that the third term in scalar potential, eq. (2.4), will be deformed by multiplying the phase factor $\exp(-2i\pi\gamma(Q^I \times Q^J + Q^I \times Q^K + Q^K \times Q^I))$. Here I, J, K take value of the integral number 1, 2, 3, 4, and we define:

$$Q^I \times Q^J \equiv Q_1^I Q_2^J - Q_1^J Q_2^I. \quad (2.11)$$

Where Q^I are the $U(1)$ charges of the fields.

[†]We denote the deformation parameter as γ to stress that it is *real*, However the supersymmetric one-parameter deformation is still called β -deformation and the non-supersymmetric three-parameter deformation to be introduced in section 6 will be called γ -deformation. We hope this will not produce confusions for the readers.

Table 2. The non-vanishing phase factors for the third term of V_B

I	J	K	phase factor
1	2	3	$i\pi\gamma$
1	2	4	$-i\pi\gamma$
1	3	2	$-i\pi\gamma$
1	3	4	$-i\pi\gamma$
1	4	2	$i\pi\gamma$
1	4	3	$i\pi\gamma$

Table 3. The non-vanishing phase factors for the second term of V_F

I	J	phase factor
1	3	$-\frac{1}{2}i\pi\gamma$
1	4	$\frac{1}{2}i\pi\gamma$
2	3	$\frac{1}{2}i\pi\gamma$
2	4	$-\frac{1}{2}i\pi\gamma$
3	1	$\frac{1}{2}i\pi\gamma$
3	2	$-\frac{1}{2}i\pi\gamma$
4	1	$-\frac{1}{2}i\pi\gamma$
4	2	$\frac{1}{2}i\pi\gamma$

So the third term now becomes:

$$\begin{aligned}
& V_{B, 3rd}^{\text{deformed}} \\
&= -\frac{1}{3} \left(\frac{2\pi}{k} \right)^2 \left[4 \sum_{\text{two of I, J, K are the same}} \overline{\text{Tr}}(Y_I^+ Y^J Y_k^+ Y^I Y_J^+ Y^K) \right. \\
&+ 4 \sum_{\text{all of I, J, K belong to (12) or (34)}} \overline{\text{Tr}}(Y_I^+ Y^J Y_k^+ Y^I Y_J^+ Y^K) \\
&+ 4 \sum_{(IJK)=(123),(143),(142)} e^{i\pi\gamma} \overline{\text{Tr}}(Y_I^+ Y^J Y_k^+ Y^I Y_J^+ Y^K) + \text{cyclic permutations} \\
&\left. + 4 \sum_{(IJK)=(132),(134),(124)} e^{-i\pi\gamma} \overline{\text{Tr}}(Y_I^+ Y^J Y_k^+ Y^I Y_J^+ Y^K) + \text{cyclic permutations} \right] .(2.12)
\end{aligned}$$

The non-vanishing phase factors in the third term of the potential V_B are listed in table 2. The other three terms in eq. (2.4) are untouched by the β -deformation.

Now we turn to consider the interaction terms between the fermions and scalars in

Table 4. The non-vanishing phase factors for the third term of V_F

I	J	K	L	phase factor
1	3	2	4	$\frac{1}{2}i\pi\gamma$
1	4	2	3	$-\frac{1}{2}i\pi\gamma$
2	3	1	4	$-\frac{1}{2}i\pi\gamma$
2	4	1	3	$\frac{1}{2}i\pi\gamma$
3	1	4	2	$-\frac{1}{2}i\pi\gamma$
3	2	4	1	$\frac{1}{2}i\pi\gamma$
4	1	3	2	$\frac{1}{2}i\pi\gamma$
4	2	3	1	$-\frac{1}{2}i\pi\gamma$

eq. (2.5). After the deformation, these terms become

$$\begin{aligned}
V_F^{\text{deformed}} = & \frac{2\pi i}{k} \text{Tr} \left[Y_I^+ Y^I \Psi^{+J} \Psi_J - 2 \sum_{I=J, \text{ or } I, J \in \{1,2\}, \text{ or } I, J \in \{3,4\}} Y_I^+ Y^J \Psi^{+I} \Psi_J \right. \\
& - 2 \sum_{(I,J)=(1,4),(2,3),(3,1),(4,2)} e^{\frac{i}{2}\pi\gamma} Y_I^+ Y^J \Psi^{+I} \Psi_J \\
& - 2 \sum_{(I,J)=(1,3),(2,4),(3,2),(4,1)} e^{-\frac{i}{2}\pi\gamma} Y_I^+ Y^J \Psi^{+I} \Psi_J \\
& + \sum_{(IJKL)=(1324),(2413),(3241),(4132)} e^{\frac{i}{2}\pi i\gamma} Y_I^+ \Psi_J Y_K^+ \Psi_L \\
& + \sum_{(IJKL)=(1423),(2314),(3142),(4231)} e^{-\frac{i}{2}\pi i\gamma} Y_I^+ \Psi_J Y_K^+ \Psi_L \\
& \left. + \sum_{\text{other terms}} \epsilon^{IJKL} \Psi_I^+ \Psi_J Y_K^+ \Psi_L \right] + \text{h.c} \tag{2.13}
\end{aligned}$$

We list the non-vanishing phase factors multiplying the second and the third terms of V_F get changed in tables 3 and 4.

3 Two-loop anomalous dimensions in the scalar sector

Now we compute the two-loop planar contributions to the anomalous dimensions for the composite local operators in the scalar sector:

$$\mathcal{O}_{J_1 \dots J_L}^{I_1 \dots I_L} \equiv \text{Tr} \left(Y^{I_1} Y_{J_1}^\dagger Y^{I_2} Y_{J_2}^\dagger \dots Y^{I_L} Y_{J_L}^\dagger \right), \quad L \geq 2. \tag{3.1}$$

The anomalous dimension matrix can be expressed as Hamiltonian acting on an alternating $SU(4)$ spin chain with the spins at odd lattice sides in the fundamental representation (**4**) and the spins at even lattices in the anti-fundamental representation ($\bar{\mathbf{4}}$). The length of the spin chain is $2L$. The involved Feynman diagrams are the same as the ones in [15, 16].




Figure 1. The contribution to anomalous dimension of \mathcal{O} from two loop contribution from scalar sextet interaction. In this context, the horizontal lines represent the operators and the ordered vertical lines denote the contraction between the two operators of the fields included in trace.




Figure 2. The contribution to anomalous dimension of \mathcal{O} from two loop contribution of gauge and fermion exchange interaction. The internal waved lines and dashed line stand gluon and scalar respectively.

Since only the third terms in V_B is modified by the deformation, one can see by some computations based on the Feynman diagram in Fig. 1 that the Hamiltonian from V_B ,

$$H_B = \frac{\lambda^2}{2} \sum_{i=1}^{2L} [\mathbb{I} - 2\mathbb{P}_{i,i+2} - \mathbb{K}_{i,i+1} + \mathbb{P}_{i,i+2}\mathbb{K}_{i,i+1} + \mathbb{K}_{i,i+1}\mathbb{P}_{i,i+1}], \quad (3.2)$$

is now changed into

$$\tilde{H}_B = \frac{\lambda^2}{2} \sum_{i=1}^{2L} [\mathbb{I} - 2\tilde{\mathbb{P}}_{i,i+2} - \mathbb{K}_{i,i+1} + \mathbb{P}_{i,i+2}\mathbb{K}_{i,i+1} + \mathbb{K}_{i,i+1}\mathbb{P}_{i,i+1}], \quad (3.3)$$

where $\lambda \equiv N/k$ is the 't Hooft coupling of ABJM theory, and definition of $\mathbb{I}, \mathbb{P}, \mathbb{K}$ are

$$\mathbb{I}_{KL}^{IJ} = \delta_K^I \delta_L^J, \quad \mathbb{P}_{KL}^{IJ} = \delta_L^I \delta_K^J, \quad \mathbb{K}_{KL}^{IJ} = \delta^{IJ} \delta_{KL}. \quad (3.4)$$

The definition of $\tilde{\mathbb{P}}_{i,i+2}$ is

$$\begin{aligned} \left(\tilde{\mathbb{P}}_{i,i+2} \right)_{J_i J_{i+1} J_{i+2}}^{I_i I_{i+1} I_{i+2}} &\equiv \exp(-i\pi\gamma(Q^{J_i} \times Q^{J_{i+1}} + Q^{J_{i+1}} \times Q^{J_{i+2}} + Q^{J_{i+2}} \times Q^{J_i})) \\ &\times (\mathbb{P}_{i,i+2})_{J_i J_{i+1} J_{i+2}}^{I_i I_{i+1} I_{i+2}}. \end{aligned} \quad (3.5)$$

The contributions to the anomalous dimension of operator (3.1) from gauge and fermion exchange interaction are relevant to Feynman diagrams in Fig. 2. The wave




Figure 3. The contribution to wave function renormalization of Y, Y^+ from two loop contribution of diamagnetic gauge interactions.




Figure 4. The contribution to wave function renormalization of Y, Y^+ is from two loop contribution of paramagnetic gauge interactions.




Figure 5. The contribution to wave function renormalization of Y, Y^+ is from two loop contribution of Chern-Simons interaction.




Figure 6. The contribution to wave function renormalization of Y, Y^+ is from two loop contribution of fermion pair interaction to wave function renormalization.

function renormalization of Y, Y^+ will also make contributions to anomalous dimension of composite operators in eq. (3.1). There are three kinds of nonzero contribution arise from interactions involving gauge boson loops, vertices given in V_F^{deformed} and gauge-matter




Figure 7. The contribution to wave function renormalization of Y , Y^+ is from two loop contribution of vacuum polarization.

interactions. The relevant Feynman diagrams are given in Fig. 3-Fig. 5, Fig. 6 and Fig. 7, respectively. We find that these contributions

$$H_F = \lambda^2 \sum_{i=1}^{2L} \mathbb{K}_{i,i+1}, \quad (3.6)$$

$$H_{\text{gauge}} = \lambda^2 \sum_{i=1}^{2L} \left(-\frac{1}{4} \mathbb{I} - \frac{1}{2} \mathbb{K}_{i,i+1} \right), \quad (3.7)$$

$$H_Z = \lambda^2 \sum_{i=1}^{2L} \frac{3}{4} \mathbb{I}, \quad (3.8)$$

are the same as the undeformed case.

By summing over all of these contributions, we get [‡]

$$\begin{aligned} \tilde{H}_{\text{total}} &= \tilde{H}_B + H_F + H_{\text{gauge}} + H_Z \\ &= \lambda^2 \sum_{i=1}^{2L} \left(\mathbb{I} - \tilde{\mathbb{P}}_{i,i+2} + \frac{1}{2} \mathbb{P}_{i,i+2} \mathbb{K}_{i,i+1} + \frac{1}{2} \mathbb{P}_{i,i+2} \mathbb{K}_{i+1,i+2} \right). \end{aligned} \quad (3.10)$$

We notice that, as the undeformed case [17], the computations in the ABJ theory [14] with gauge group $U(N)_k \times U(M)_{-k}$ is almost the same besides replacing the factor λ^2 in the Hamiltonian by the factor $\lambda \tilde{\lambda}$, where $\tilde{\lambda}$ is defined to be $\tilde{\lambda} \equiv M/k$. So the computations and discussions in the following applied to the scalar section in ABJ theory at two-loop level as well.

4 The R matrices of deformed spin chain

In this section, we will show that Hamiltonian obtained in the previous section is integrable by constructing the R matrix which satisfies the Yang-Baxter equation (YBE) and gives this Hamiltonian through the transfer matrix by the standard procedure.

[‡] Here our convention is

$$\begin{aligned} & (\mathbb{P}_{i,i+2} \mathbb{K}_{i+1,i+2})_{J_i J_{i+1} J_{i+2}}^{K_i K_{i+1} K_{i+2}} \\ &= (\mathbb{P}_{i,i+2})_{L_i L_{i+1} L_{i+2}}^{K_i K_{i+1} K_{i+2}} (\mathbb{K}_{i+1,i+2})_{J_i J_{i+1} J_{i+2}}^{L_i L_{i+1} L_{i+2}} \end{aligned} \quad (3.9)$$

For the alternating spin chain, we need four R-matrices $\widetilde{\mathfrak{R}}_{ij}^{44}(u)$, $\widetilde{\mathfrak{R}}_{ij}^{4\bar{4}}(u)$, $\widetilde{\mathfrak{R}}_{ij}^{\bar{4}4}(u)$, $\widetilde{\mathfrak{R}}_{ij}^{\bar{4}\bar{4}}(u)$ acting on the space $V_i \otimes V_j$. Here the upper indices of $\widetilde{\mathfrak{R}}_{ij}^{44}(u)$ denote $SU(4)$ representations related to the two spaces and u denotes spectral parameter. We defined these R-matrices as

$$\widetilde{\mathfrak{R}}^{44}(u)_{KL}^{IJ} = \exp(-\frac{i}{2}\pi\gamma(Q^J \times Q^I - Q^K \times Q^L))\mathfrak{R}^{44}(u)_{KL}^{IJ}, \quad (4.1)$$

$$\widetilde{\mathfrak{R}}^{4\bar{4}}(u)_{KL}^{IJ} = \exp(\frac{i}{2}\pi\gamma(Q^J \times Q^I - Q^K \times Q^L))\mathfrak{R}^{4\bar{4}}(u)_{KL}^{IJ}, \quad (4.2)$$

$$\widetilde{\mathfrak{R}}^{\bar{4}4}(u)_{KL}^{IJ} = \exp(\frac{i}{2}\pi\gamma(Q^J \times Q^I - Q^K \times Q^L))\mathfrak{R}^{\bar{4}4}(u)_{KL}^{IJ}, \quad (4.3)$$

$$\widetilde{\mathfrak{R}}^{\bar{4}\bar{4}}(u)_{KL}^{IJ} = \exp(-\frac{i}{2}\pi\gamma(Q^J \times Q^I - Q^K \times Q^L))\mathfrak{R}^{\bar{4}\bar{4}}(u)_{KL}^{IJ}, \quad (4.4)$$

where

$$\mathfrak{R}^{44}(u) = u\mathbb{I} + \mathbb{P}, \quad (4.5)$$

$$\mathfrak{R}^{4\bar{4}}(u) = -(u+2)\mathbb{I} + \mathbb{K}, \quad (4.6)$$

$$\mathfrak{R}^{\bar{4}4}(u) = -(u+2)\mathbb{I} + \mathbb{K}, \quad (4.7)$$

$$\mathfrak{R}^{\bar{4}\bar{4}}(u) = u\mathbb{I} + \mathbb{P}, \quad (4.8)$$

are the R-matrices before deformation [15, 16]. From above formulas, we can get:

$$\widetilde{\mathfrak{R}}^{44}(u)_{KL}^{IJ} = u \exp(-i\pi\gamma Q^J \times Q^I) \delta_K^I \delta_L^J + \delta_L^I \delta_K^J, \quad (4.9)$$

$$\widetilde{\mathfrak{R}}^{4\bar{4}}(u)_{KL}^{IJ} = -(u+2) \exp(i\pi\gamma Q^J \times Q^I) \delta_K^I \delta_L^J + \delta^{IJ} \delta_{KL}, \quad (4.10)$$

$$\widetilde{\mathfrak{R}}^{\bar{4}4}(u)_{KL}^{IJ} = -(u+2) \exp(i\pi\gamma Q^J \times Q^I) \delta_K^I \delta_L^J + \delta^{IJ} \delta_{KL}, \quad (4.11)$$

$$\widetilde{\mathfrak{R}}^{\bar{4}\bar{4}}(u)_{KL}^{IJ} = u \exp(-i\pi\gamma Q^J \times Q^I) \delta_K^I \delta_L^J + \delta_L^I \delta_K^J. \quad (4.12)$$

The R matrices before deformation satisfy YBE [15, 16]:

$$\mathfrak{R}^{44}_{12}(u-v)\mathfrak{R}^{44}_{13}(u)\mathfrak{R}^{44}_{23}(v) = \mathfrak{R}^{44}_{23}(v)\mathfrak{R}^{44}_{13}(u)\mathfrak{R}^{44}_{12}(u-v), \quad (4.13)$$

$$\mathfrak{R}^{44}_{12}(u-v)\mathfrak{R}^{4\bar{4}}_{13}(u)\mathfrak{R}^{4\bar{4}}_{23}(v) = \mathfrak{R}^{4\bar{4}}_{23}(v)\mathfrak{R}^{4\bar{4}}_{13}(u)\mathfrak{R}^{44}_{12}(u-v), \quad (4.14)$$

$$\mathfrak{R}^{\bar{4}4}_{12}(u-v)\mathfrak{R}^{\bar{4}4}_{13}(u)\mathfrak{R}^{\bar{4}4}_{23}(v) = \mathfrak{R}^{\bar{4}4}_{23}(v)\mathfrak{R}^{\bar{4}4}_{13}(u)\mathfrak{R}^{\bar{4}4}_{12}(u-v), \quad (4.15)$$

$$\mathfrak{R}^{\bar{4}\bar{4}}_{12}(u-v)\mathfrak{R}^{\bar{4}\bar{4}}_{13}(u)\mathfrak{R}^{\bar{4}\bar{4}}_{23}(v) = \mathfrak{R}^{\bar{4}\bar{4}}_{23}(v)\mathfrak{R}^{\bar{4}\bar{4}}_{13}(u)\mathfrak{R}^{\bar{4}\bar{4}}_{12}(u-v). \quad (4.16)$$

As in [50], the choice of the phases in eqs. (4.1-4.4) are such that the R matrices after deformation still satisfy the YBE:

$$\widetilde{\mathfrak{R}}^{44}_{12}(u-v)\widetilde{\mathfrak{R}}^{44}_{13}(u)\widetilde{\mathfrak{R}}^{44}_{23}(v) = \widetilde{\mathfrak{R}}^{44}_{23}(v)\widetilde{\mathfrak{R}}^{44}_{13}(u)\widetilde{\mathfrak{R}}^{44}_{12}(u-v), \quad (4.17)$$

$$\widetilde{\mathfrak{R}}^{44}_{12}(u-v)\widetilde{\mathfrak{R}}^{4\bar{4}}_{13}(u)\widetilde{\mathfrak{R}}^{4\bar{4}}_{23}(v) = \widetilde{\mathfrak{R}}^{4\bar{4}}_{23}(v)\widetilde{\mathfrak{R}}^{4\bar{4}}_{13}(u)\widetilde{\mathfrak{R}}^{44}_{12}(u-v), \quad (4.18)$$

$$\widetilde{\mathfrak{R}}^{\bar{4}4}_{12}(u-v)\widetilde{\mathfrak{R}}^{\bar{4}4}_{13}(u)\widetilde{\mathfrak{R}}^{\bar{4}4}_{23}(v) = \widetilde{\mathfrak{R}}^{\bar{4}4}_{23}(v)\widetilde{\mathfrak{R}}^{\bar{4}4}_{13}(u)\widetilde{\mathfrak{R}}^{\bar{4}4}_{12}(u-v), \quad (4.19)$$

$$\widetilde{\mathfrak{R}}^{\bar{4}\bar{4}}_{12}(u-v)\widetilde{\mathfrak{R}}^{\bar{4}\bar{4}}_{13}(u)\widetilde{\mathfrak{R}}^{\bar{4}\bar{4}}_{23}(v) = \widetilde{\mathfrak{R}}^{\bar{4}\bar{4}}_{23}(v)\widetilde{\mathfrak{R}}^{\bar{4}\bar{4}}_{13}(u)\widetilde{\mathfrak{R}}^{\bar{4}\bar{4}}_{12}(u-v). \quad (4.20)$$

By introducing an auxiliary space V_0 , we can define the following two transfer T-matrices [§]:

$$\begin{aligned} \widetilde{T}_0(u, a) = & 2^{-L} \widetilde{\mathfrak{R}}^{44}_{01}(u) \widetilde{\mathfrak{R}}^{44}_{02}(u+a) \widetilde{\mathfrak{R}}^{44}_{03}(u) \widetilde{\mathfrak{R}}^{44}_{04}(u+a) \cdots \\ & \widetilde{\mathfrak{R}}^{44}_{0(2L-1)}(u) \widetilde{\mathfrak{R}}^{44}_{0(2L)}(u+a), \end{aligned} \quad (4.21)$$

$$\begin{aligned} \widetilde{\bar{T}}_0(u, \bar{a}) = & 2^{-L} \widetilde{\mathfrak{R}}^{44}_{01}(u+\bar{a}) \widetilde{\mathfrak{R}}^{44}_{02}(u) \widetilde{\mathfrak{R}}^{44}_{03}(u+\bar{a}) \widetilde{\mathfrak{R}}^{44}_{04}(u) \cdots \\ & \widetilde{\mathfrak{R}}^{44}_{0(2L-1)}(u+\bar{a}) \widetilde{\mathfrak{R}}^{44}_{0(2L)}(u). \end{aligned} \quad (4.22)$$

The YBE will lead to the following relations:

$$\widetilde{\mathfrak{R}}^{44}_{00'}(\nu - \mu) \widetilde{T}_0(\nu, a) \widetilde{T}_{0'}(\mu, a) = \widetilde{T}_{0'}(\mu, a) \widetilde{T}_0(\nu, a) \widetilde{\mathfrak{R}}^{44}_{00'}(\nu - \mu), \quad (4.23)$$

$$\widetilde{\mathfrak{R}}^{44}_{00'}(\nu - \mu) \widetilde{\bar{T}}_0(\nu, \bar{a}) \widetilde{\bar{T}}_{0'}(\mu, \bar{a}) = \widetilde{\bar{T}}_{0'}(\mu, \bar{a}) \widetilde{\bar{T}}_0(\nu, \bar{a}) \widetilde{\mathfrak{R}}^{44}_{00'}(\nu - \mu), \quad (4.24)$$

$$\widetilde{\mathfrak{R}}^{44}_{00'}(\nu - \mu + a) \widetilde{T}_0(\nu, a) \widetilde{\bar{T}}_{0'}(\mu, -a) = \widetilde{\bar{T}}_{0'}(\mu, -a) \widetilde{T}_0(\nu, a) \widetilde{\mathfrak{R}}^{44}_{00'}(\nu - \mu + a). \quad (4.25)$$

Then the traces of the two T-matrices

$$\widetilde{\tau}(\nu, a) = \text{Tr}_0 \widetilde{T}_0(\nu, a), \quad (4.26)$$

$$\widetilde{\bar{\tau}}(\nu, a) = \text{Tr}_0 \widetilde{\bar{T}}_0(\nu, a), \quad (4.27)$$

satisfy

$$\begin{aligned} [\widetilde{\tau}(\nu, a), \widetilde{\tau}(\mu, a)] &= 0 \\ [\widetilde{\bar{\tau}}(\nu, \bar{a}), \widetilde{\bar{\tau}}(\mu, \bar{a})] &= 0, \\ [\widetilde{\tau}(\nu, a), \widetilde{\bar{\tau}}(\mu, -a)] &= 0. \end{aligned} \quad (4.28)$$

From now on, we restrict a to be purely imaginary so that $\widetilde{\tau}(\nu, a)$ and $\widetilde{\bar{\tau}}(\mu, \bar{a})$ commute with each other.

Now we compute the Hamiltonian from the R-matrices:

$$H = \frac{\partial \log \widetilde{\tau}(u, a)}{\partial u} \Big|_{u=0} + \frac{\partial \log \widetilde{\bar{\tau}}(u, \bar{a})}{\partial u} \Big|_{u=0}. \quad (4.29)$$

After some computations, we get

$$H = \sum_{i=1}^{2L} H_i, \quad (4.30)$$

with

$$\begin{aligned} H_{2l-1} = & \frac{1}{a^2 - 4} ((a-2)\mathbb{I} + (a^2 - 4)\widetilde{\mathbb{P}}_{2l-1, 2l+1} \\ & - (a-2)\mathbb{P}_{2l-1, 2l+1}\mathbb{K}_{2l-1, 2l} + (a+2)\mathbb{P}_{2l-1, 2l+1}\mathbb{K}_{2l, 2l+1}), \end{aligned} \quad (4.31)$$

$$\begin{aligned} H_{2l} = & \frac{1}{a^2 - 4} (-(a+2)\mathbb{I} + (a^2 - 4)\widetilde{\mathbb{P}}_{2l, 2l+2} \\ & + (a+2)\mathbb{P}_{2l, 2l+2}\mathbb{K}_{2l, 2l+1} - (a-2)\mathbb{P}_{2l, 2l+2}\mathbb{K}_{2l+1, 2l+2}). \end{aligned} \quad (4.32)$$

[§]Comparing with the T-matrices in [16], we include the constant factor 2^{-L} as in [15].

Here we have already used the relation $\bar{a} = -a$. The details of the computations are deferred to the Appendix. After setting $a = 0$, multiplying H_i by -1 , then shifting H_i by $\frac{3}{2}\mathbb{I}$, the above Hamiltonian coincides with the one in the previous section obtained from the perturbative computations in field theory side.

5 Eigenvalues of deformed spin chain Hamiltonian and Bethe ansatz equations

In previous section, we constructed transfer matrix and Hamiltonian of the deformed spin chain. We now derive the Bethe ansatz equation through diagonalizing the transfer matrices. By choosing the ground state or highest-weight state as $|\bar{1}\bar{4}1\bar{4}\cdots\rangle$ and introducing three sets of Bethe roots $(l_a, m_b, r_c), 1 \leq a \leq N_l, 1 \leq b \leq N_m, 1 \leq c \leq N_r$, we can get the eigenvalues of $\tilde{\tau}(\nu, 0)$ as

$$\begin{aligned} \tilde{\Lambda}(\nu) = & 2^{-L}(\nu+1)^L(-\nu-2)^L \exp\left(\frac{i}{4}\pi\gamma L + \frac{i}{4}\gamma N_m - \frac{i}{2}\pi\gamma N_r\right) \prod_{a=1}^{N_l} \frac{\nu + il_a - \frac{1}{2}}{\nu + il_a + \frac{1}{2}} \\ & + 2^{-L}(\nu+1)^L(-\nu)^L \exp\left(\frac{i}{4}\pi\gamma L + \frac{i}{4}\gamma N_m - \frac{i}{2}\pi\gamma N_l\right) \prod_{c=1}^{N_r} \frac{\nu + ir_c + \frac{5}{2}}{\nu + ir_c + \frac{3}{2}} \\ & + 2^{-L}(-\nu-2)^L\nu^L \exp\left(-\frac{i}{4}\pi\gamma L - \frac{i}{4}\pi\gamma N_m + \frac{i}{2}\pi\gamma N_r\right) \prod_{a=1}^{N_l} \frac{\nu + il_a + \frac{3}{2}}{\nu + il_a + \frac{1}{2}} \prod_{b=1}^{N_m} \frac{\nu + im_b}{\nu + im_b + 1} \\ & + 2^{-L}(-\nu-2)^L\nu^L \exp\left(-\frac{i}{4}\pi\gamma L - \frac{i}{4}\pi\gamma N_m + \frac{i}{2}\pi\gamma N_l\right) \prod_{c=1}^{N_r} \frac{\nu + ir_c + \frac{1}{2}}{\nu + ir_c + \frac{3}{2}} \prod_{b=1}^{N_m} \frac{\nu + im_b + 2}{\nu + im_b + 1}. \end{aligned} \quad (5.1)$$

As the undeformed case [15, 16], N_l, N_m, N_r should satisfy:

$$2N_l \leq L + N_m, 2N_r \leq L + N_m, 2N_m \leq N_l + N_r. \quad (5.2)$$

Similarly, we can get the eigenvalues of $\tilde{\tilde{\tau}}(\nu, 0)$ as

$$\begin{aligned} \tilde{\tilde{\Lambda}}(\nu) = & 2^{-L}(-\nu)^L(\nu+1)^L \exp\left(-\frac{i}{4}\pi\gamma L - \frac{i}{4}\pi\gamma N_m + \frac{i}{2}\pi\gamma N_r\right) \prod_{a=1}^{N_l} \frac{\nu + il_a + \frac{5}{2}}{\nu + il_a + \frac{3}{2}} \\ & + 2^{-L}(\nu+1)^L(-\nu-2)^L \exp\left(-\frac{i}{4}\pi\gamma L - \frac{i}{4}\pi\gamma N_m + \frac{i}{2}\pi\gamma N_l\right) \prod_{c=1}^{N_r} \frac{\nu + ir_c - \frac{1}{2}}{\nu + ir_c + \frac{1}{2}} \\ & + 2^{-L}(-\nu-2)^L\nu^L \exp\left(\frac{i}{4}\pi\gamma L + \frac{i}{4}\pi\gamma N_m - \frac{i}{2}\pi\gamma N_r\right) \prod_{a=1}^{N_l} \frac{\nu + il_a + \frac{1}{2}}{\nu + il_a + \frac{3}{2}} \prod_{b=1}^{N_m} \frac{\nu + im_b + 2}{\nu + im_b + 1} \\ & + 2^{-L}(-\nu-2)^L\nu^L \exp\left(\frac{i}{4}\pi\gamma L + \frac{i}{4}\pi\gamma N_m - \frac{i}{2}\pi\gamma N_l\right) \prod_{b=1}^{N_m} \frac{\nu + im_b}{\nu + im_b + 1} \prod_{c=1}^{N_r} \frac{\nu + ir_c + \frac{3}{2}}{\nu + ir_c + \frac{1}{2}}. \end{aligned} \quad (5.3)$$

By demanding the residue vanishes at every pole of $\tilde{\Lambda}(\nu)$, we get the following set of Bethe Ansatz equations.

$$\begin{aligned} \exp\left(-\frac{i}{2}\pi\gamma L - \frac{i}{2}\pi\gamma N_m + i\pi\gamma N_l\right) \left(\frac{l_a - \frac{i}{2}}{l_a + \frac{i}{2}}\right)^L &= \prod_{a' \neq a} \frac{l_a - l_{a'} - i}{l_a - l_{a'} + i} \prod_{b=1}^{N_m} \frac{l_a - m_b + \frac{i}{2}}{l_a - m_b - \frac{i}{2}}, \\ \exp\left(\frac{i}{2}\pi\gamma L + \frac{i}{2}\pi\gamma N_m - i\pi\gamma N_l\right) \left(\frac{r_c - \frac{i}{2}}{r_c + \frac{i}{2}}\right)^L &= \prod_{b=1}^{N_m} \frac{r_c - m_b + \frac{i}{2}}{r_c - m_b - \frac{i}{2}} \prod_{c' \neq c} \frac{r_c - r_{c'} - i}{r_c - r_{c'} + i}, \\ \exp\left(\frac{i}{2}\pi\gamma N_l - \frac{i}{2}\pi\gamma N_r\right) &= \prod_{a=1}^{N_l} \frac{m_b - l_a - \frac{i}{2}}{m_b - l_a + \frac{i}{2}} \prod_{b \neq b'} \frac{m_b - m_{b'} - i}{m_b - m_{b'} + i} \prod_{c=1}^{N_r} \frac{m_b - r_c + \frac{i}{2}}{m_b - r_c - \frac{i}{2}}. \end{aligned} \quad (5.4)$$

We will get the same set of equations if we start with $\tilde{\tilde{\Lambda}}(\nu)$ instead. It is a consistent check that the same set of Bethe ansatz equations remove potential simple pole terms for $\tilde{\Lambda}(\nu)$ and $\tilde{\tilde{\Lambda}}(\nu)$. One can see the equations just above go back to Bethe ansatz equations given in [15, 16] by setting $\gamma = 0$.

From the above eigenvalues, we can get the total momentum as

$$\begin{aligned} P_{total} &= \frac{1}{i} \left[\log \tilde{\Lambda}(0) + \log \tilde{\tilde{\Lambda}}(0) \right] \\ &= \frac{1}{i} \left[-\frac{i}{2}\pi\gamma N_r + \frac{i}{2}\pi\gamma N_l + \sum_{a=1}^{N_l} \log \frac{il_a - \frac{1}{2}}{il_a + \frac{1}{2}} + \sum_{c=1}^{N_r} \log \frac{ir_c - \frac{1}{2}}{ir_c + \frac{1}{2}} \right]. \end{aligned} \quad (5.5)$$

From the vanishing of the total momentum, we can obtain the following constraint:

$$1 = \exp\left(-\frac{i}{2}\pi\gamma N_r + \frac{i}{2}\pi\gamma N_l\right) \prod_{a=1}^{N_l} \frac{il_a - \frac{1}{2}}{il_a + \frac{1}{2}} \prod_{c=1}^{N_r} \frac{ir_c - \frac{1}{2}}{ir_c + \frac{1}{2}}. \quad (5.6)$$

The total energy is

$$\begin{aligned} E_{total} &= - \left[\frac{d}{d\nu} \log \tilde{\Lambda}(\nu) + \frac{d}{d\nu} \log \tilde{\tilde{\Lambda}}(\nu) \right] \Big|_{\nu=0} \\ &= \left(\sum_{a=1}^{N_l} \frac{1}{l_a^2 + \frac{1}{4}} + \sum_{c=1}^{N_r} \frac{1}{r_c^2 + \frac{1}{4}} \right). \end{aligned} \quad (5.7)$$

From the total energy, we can get the eigenvalues of the anomalous dimension matrix of the operators in the scalar sector eq. (3.1) in the β -deformed ABJM theory.

6 The non-supersymmetric three-parameter γ -deformation of ABJ(M) theory

The three-parameter deformation can be performed by replacing all of the ordinary product fg of two fields f and g in the Lagrangian by the following star product [74]:

$$f * g = e^{i\pi\gamma_i Q_j^f Q_k^g \epsilon^{ijk}} fg, \quad (6.1)$$

Table 5. $U(1)^3$ charges of the scalars and fermions of the ABJM theory used for γ -deformation.

	Y^1	Y^2	Y^3	Y^4	$\Psi^{\dagger 1}$	$\Psi^{\dagger 2}$	$\Psi^{\dagger 3}$	$\Psi^{\dagger 4}$
$U(1)_1$	$\frac{1}{2}$	$-\frac{1}{2}$	0	0	$\frac{1}{2}$	$-\frac{1}{2}$	0	0
$U(1)_2$	0	0	$\frac{1}{2}$	$-\frac{1}{2}$	0	0	$\frac{1}{2}$	$-\frac{1}{2}$
$U(1)_3$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$

where $Q_i^f, i = 1, 2, 3$ are three global $U(1)$ charges carrying by the field f and γ_i 's are three real deformation parameters. We choose the $U(1)_i$ charges for the scalars and the fermions as in table 5. The gauge field is neutral under these symmetries. One can see that this deformation degenerates to the one of β -deformation by setting deformation parameters $\gamma_1 = \gamma_2 = 0, \gamma_3 = \gamma$.

After analogous calculation as in section 3, one can find that in the scalar sector and at the two-loop level, the Hamiltonian is still given by eq. (3.10). The differences between β - and γ -deformations may appear at higher loop orders or in other sectors of the theories.

7 Conclusion and Discussions

In this note, we start a study on the integrable structure of β - and γ -deformed ABJ(M) theory, beginning with the scalar sector at the two-loop level in the planar limit. We first perform perturbative computations of the anomalous dimension matrix and express the result as a Hamiltonian acting on alternating $SU(4)$ spin chain. We find that only one term in the Hamiltonian is deformed and that the differences between β -deformation and γ -deformation are invisible in this sector at two loop level. As the undeformed case, the difference between Hamiltonian for deformed ABJM theory and deformed ABJ theory only appears in the prefactor. So in this sector and at this order of the perturbation theory, the violation of parity invariance in the deformed ABJ theory does not affect the integrability. Based on the structure of the deformations, we choose a suitable deformation of the R-matrices. Then the Bethe ansatz equations are obtained through diagonalizing the transfer matrices.

There are several directions worth pursuing. The study here in the field theory side can be extended to full sector and/or higher loop order as in the undeformed case [79]–[85]. It is also interesting to reproduce the Bethe ansatz equation starting from the S-matrix of the spin chain based on the studies in [18, 64, 65, 72]. In the string theory side, one could try to construct the Lax pair and the infinite number of conserved currents on the worldsheet. Even for the undeformed case, the story of IIA string on $AdS_4 \times CP^3$ has already been much richer than the one of IIB string on $AdS_5 \times S^5$, partly because that now the $OSp(6|4)/U(3) \times SO(1, 3)$ coset action can only describe a subset of the complete Green-Schwarz action [21–23].

Acknowledgements

The authors are grateful to Dongsu Bak, Bin Chen, Peng Gao, Miao Li, Xiao Liu, Jian-Xin Lu, Jian-Feng Wu, Zhi-Guang Xiao, Jie Yang and Hossein Yavartanoo for useful

discussions. JW would like to thank SISSA, ICTS-USTC and School of Mathematical Sciences, USTC for warm hospitality. He is also very thankful to the organizers and participants of Symposium on Strings and Particle Physics 2012 held at CTP, SCU and International Workshop on Quantum Aspects of Black Holes held in CQeST, Sogang University. This work was supported in part by the National Natural Science Foundation of China (No.10821504 (SH), No.10975168 (SH), No.11035008 (SH), No. 11105154 (JW), and No. 11222549 (JW)), and in part by the Ministry of Science and Technology of China under Grant No. 2010CB833004. SH also would like to appreciate the general financial support from China Postdoctoral Science Foundation No. 2012M510562. JW gratefully acknowledges the support of K. C. Wong Education Foundation and Youth Innovation Promotion Association, CAS as well.

A Hamiltonian from deformed R-matrices

In this appendix, we compute the Hamiltonian of deformed spin chain in eq. (4.29). For this, we should compute $\tilde{\tau}'(0, a)$, where the prime denotes the derivative with respect to spectrum parameter u .

$$\begin{aligned} \tilde{\tau}'(0, a) = & 2^{-L} \text{Tr} \sum_i \widetilde{\mathfrak{R}}^{44}_{01}(0) \dots \frac{d\widetilde{\mathfrak{R}}^{44}_{0(2i-1)}(u)}{du} \Big|_{u=0} \dots \widetilde{\mathfrak{R}}^{44}_{0(2L)}(a) \\ & + 2^{-L} \text{Tr} \sum_i \widetilde{\mathfrak{R}}^{44}_{01}(0) \dots \frac{d\widetilde{\mathfrak{R}}^{44}_{0(2i)}(u+a)}{du} \Big|_{u=0} \dots \widetilde{\mathfrak{R}}^{44}_{0(2L)}(a). \end{aligned} \quad (\text{A.1})$$

Where the spectrum parameter u has been set as vanishing. The i -th term in first part of eq. (A.1) can be written down as following

$$\begin{aligned} & 2^{-L} (\mathbb{P}_{01})_{K_0 J_1}^{K_1 I_1} \widetilde{\mathfrak{R}}^{44}_{02}(a)_{K_1 J_2}^{K_2 I_2} \dots \widetilde{\mathfrak{R}}^{44}_{0(2i-2)}(a)_{K_{2i-3} J_{2i-2}}^{K_{2i-2} I_{2i-2}} \delta_{K_{2i-2}}^{K_{2i-1}} e^{i\pi\gamma Q^{K_{2i-1} \times Q^{I_{2i-1}}}} \\ & \delta_{J_{2i-1}}^{I_{2i-1}} \widetilde{\mathfrak{R}}^{44}_{0(2i)}(a)_{K_{2i-1} J_{2i}}^{K_{2i} I_{2i}} \dots (\mathbb{P}_{0(2L-1)})_{K_{2L-2} J_{2L-1}}^{K_{2L-1} I_{2L-1}} \widetilde{\mathfrak{R}}^{44}_{0(2L)}(a)_{K_{2L-1} J_{2L}}^{K_0 I_{2L}} \\ = & 2^{-L} \widetilde{\mathfrak{R}}^{44}_{02}(a)_{J_1 J_2}^{I_3 I_2} \dots \widetilde{\mathfrak{R}}^{44}_{0(2i-2)}(a)_{J_{2i-3} J_{2i-2}}^{K_{2i-1} I_{2i-2}} e^{i\pi\gamma Q^{K_{2i-1} \times Q^{I_{2i-1}}}} \delta_{J_{2i-1}}^{I_{2i-1}} \widetilde{\mathfrak{R}}^{44}_{0(2i)}(a)_{K_{2i-1} J_{2i}}^{I_{2i+1} I_{2i}} \\ & \dots \widetilde{\mathfrak{R}}^{44}_{0(2L)}(a)_{J_{2L-1} J_{2L}}^{I_1 I_{2L}}. \end{aligned} \quad (\text{A.2})$$

The i -th term in the second part of eq. (A.1) is

$$\begin{aligned} & 2^{-L} (\mathbb{P}_{01})_{K_0 J_1}^{K_1 I_1} \widetilde{\mathfrak{R}}^{44}_{02}(a)_{K_1 J_2}^{K_2 I_2} \dots \widetilde{\mathfrak{R}}^{44}_{0(2i-2)}(a)_{K_{2i-3} J_{2i-2}}^{K_{2i-2} I_{2i-2}} (\mathbb{P}_{0(2i-1)})_{K_{2i-2} J_{2i-1}}^{K_{2i-1} I_{2i-1}} (-\mathbb{I})_{K_{2i-1} J_{2i}}^{K_{2i} I_{2i}} \\ & e^{-i\pi\gamma Q^{K_{2i} \times Q^{I_{2i}}}} (\mathbb{P}_{0(2i+1)})_{K_{2i} J_{2i+1}}^{K_{2i+1} I_{2i+1}} \widetilde{\mathfrak{R}}^{44}_{0(2i+2)}(a)_{K_{2i+1} J_{2i+2}}^{K_{2i+2} I_{2i+2}} \dots \widetilde{\mathfrak{R}}^{44}_{0(2L)}(a)_{K_{2L-1} J_{2L}}^{K_0 I_{2L}} \\ = & -2^{-L} \widetilde{\mathfrak{R}}^{44}_{02}(a)_{J_1 J_2}^{I_3 I_2} \dots \widetilde{\mathfrak{R}}^{44}_{0(2i-2)}(a)_{J_{2i-3} J_{2i-2}}^{I_{2i-1} I_{2i-2}} \delta_{J_{2i}}^{I_{2i}} \delta_{J_{2i-1}}^{I_{2i+1}} e^{-i\pi\gamma Q^{I_{2i+1} \times Q^{I_{2i}}}} \\ & \widetilde{\mathfrak{R}}^{44}_{0(2i+2)}(a)_{J_{2i+1} J_{2i+2}}^{I_{2i+3} I_{2i+2}} \dots \widetilde{\mathfrak{R}}^{44}_{0(2L)}(a)_{J_{2L-1} J_{2L}}^{I_1 I_{2L}}. \end{aligned} \quad (\text{A.3})$$

The deformed $\tilde{\tau}^{-1}(0, a)$ is

$$\tilde{\tau}^{-1}(0, a) = 2^L \left[\widetilde{\mathfrak{R}}^{44}_{0(2L)}(a) \right]_{I_1 I_{2L}}^{-1} \dots \left[\widetilde{\mathfrak{R}}^{44}_{0(2)}(a) \right]_{I_3 I_2}^{-1} \left[\widetilde{\mathfrak{R}}^{44}_{0(2)}(a) \right]_{J_1 J_2}, \quad (\text{A.4})$$

where

$$\left[\widetilde{\mathfrak{R}}^{4\bar{4}}_{0(2i)}^{-1}(a) \right]_{I_{2i+1}I_{2i}}^{J_{2i-1}J_{2i}} = - \left(\frac{1}{a+2} \mathbb{I}_{I_{2i+1}I_{2i}}^{J_{2i-1}J_{2i}} e^{i\pi\gamma Q^{J_{2i-1}} \times Q^{J_{2i}}} + \frac{1}{a^2-4} \mathbb{K}_{I_{2i+1}I_{2i}}^{J_{2i-1}J_{2i}} \right). \quad (\text{A.5})$$

One can use above formula to obtain that

$$\begin{aligned} & \widetilde{\tau}^{-1}(u, a) \widetilde{\tau}'(u, a)|_{u=0} \\ &= \sum_{i=1}^L \mathbb{I}_{J_1 \dots J_{2i-4}}^{K_1 \dots K_{2i-4}} \otimes \left[\widetilde{\mathfrak{R}}^{4\bar{4}}_{0(2i)}^{-1}(a) \right]_{I_{2i+1}I_{2i}}^{K_{2i-1}K_{2i}} \left[\widetilde{\mathfrak{R}}^{4\bar{4}}_{0(2i-2)}^{-1}(a) \right]_{I_{2i-1}I_{2i-2}}^{K_{2i-3}K_{2i-2}} \\ & \quad e^{i\pi\gamma Q^{\tilde{K}_{2i-1}} \times Q^{I_{2i-1}}} \delta_{J_{2i-1}}^{I_{2i-1}} \widetilde{\mathfrak{R}}^{4\bar{4}}_{0(2i-2)}(a)_{J_{2i-3}J_{2i-2}}^{\tilde{K}_{2i-1}I_{2i-2}} \widetilde{\mathfrak{R}}^{4\bar{4}}_{0(2i)}(a)_{\tilde{K}_{2i-1}J_{2i}}^{I_{2i+1}I_{2i}} \otimes \mathbb{I}_{J_{2i+1} \dots J_{2L}}^{K_{2i+1} \dots K_{2L}} \\ &+ \sum_{i=1}^L \mathbb{I}_{J_1 \dots J_{2i-2}}^{K_1 \dots K_{2i-2}} \otimes \left[\widetilde{\mathfrak{R}}^{4\bar{4}}_{0(2i)}^{-1}(a) \right]_{I_{2i+1}I_{2i}}^{K_{2i-1}K_{2i}} \left(-\delta_{J_{2i-1}}^{I_{2i+1}} \delta_{J_{2i}}^{I_{2i}} \right) e^{-i\pi\gamma Q^{I_{2i+1}} \times Q^{I_{2i}}} \otimes \mathbb{I}_{J_{2i+1} \dots J_{2L}}^{K_{2i+1} \dots K_{2L}} \\ &= \sum_{i=1}^L \mathbb{I}_{J_1 \dots J_{2i-4}}^{K_1 \dots K_{2i-4}} \otimes \left[\widetilde{\mathfrak{R}}^{4\bar{4}}_{0(2i)}^{-1}(a) \right]_{I_{2i+1}I_{2i}}^{K_{2i-1}K_{2i}} \left[\widetilde{\mathfrak{R}}^{4\bar{4}}_{0(2i-2)}^{-1}(a) \right]_{J_{2i-1}I_{2i-2}}^{K_{2i-3}K_{2i-2}} e^{i\pi\gamma Q^{\tilde{K}_{2i-1}} \times Q^{J_{2i-1}}} \\ & \quad \widetilde{\mathfrak{R}}^{4\bar{4}}_{0(2i-2)}(a)_{J_{2i-3}J_{2i-2}}^{\tilde{K}_{2i-1}I_{2i-2}} \widetilde{\mathfrak{R}}^{4\bar{4}}_{0(2i)}(a)_{\tilde{K}_{2i-1}J_{2i}}^{I_{2i+1}I_{2i}} \otimes \mathbb{I}_{J_{2i+1} \dots J_{2L}}^{K_{2i+1} \dots K_{2L}} \\ &- \sum_{i=1}^L \mathbb{I}_{J_1 \dots J_{2i-2}}^{K_1 \dots K_{2i-2}} \otimes \left[\widetilde{\mathfrak{R}}^{4\bar{4}}_{0(2i)}^{-1}(a) \right]_{J_{2i-1}J_{2i}}^{K_{2i-1}K_{2i}} e^{-i\pi\gamma Q^{J_{2i-1}} \times Q^{J_{2i}}} \otimes \mathbb{I}_{J_{2i+1} \dots J_{2L}}^{K_{2i+1} \dots K_{2L}} \\ &= \sum_{i=1}^L \frac{1}{a^2-4} ((a-2)\mathbb{I} + (a^2-4)\widetilde{\mathbb{P}}_{2i-1,2i+1} - (a-2)\mathbb{P}_{2i-1,2i+1}\mathbb{K}_{2i-1,2i} \\ & \quad + (a+2)\mathbb{P}_{2i-1,2i+1}\mathbb{K}_{2i,2i+1}) \end{aligned} \quad (\text{A.6})$$

Similarly, one can get:

$$\begin{aligned} & \widetilde{\tau}^{-1}(u, a) \widetilde{\tau}'(u, a)|_{u=0} \\ &= \sum_{i=1}^L \frac{1}{a^2-4} (-(a+2)\mathbb{I} + (a^2-4)\widetilde{\mathbb{P}}_{2i,2i+2} + (a+2)\mathbb{P}_{2i,2i+2}\mathbb{K}_{2i,2i+1} \\ & \quad - (a-2)\mathbb{P}_{2i,2i+2}\mathbb{K}_{2i+1,2i+2}), \end{aligned} \quad (\text{A.7})$$

where we have use the fact that a is purely imaginary. From these two equations, we can get the Hamiltonian given in the main text.

References

- [1] N. Beisert, C. Ahn, L. F. Alday, Z. Bajnok, J. M. Drummond, L. Freyhult, N. Gromov and R. A. Janik *et al.*, “Review of AdS/CFT Integrability: An Overview,” *Lett. Math. Phys.* **99**, 3 (2012) [arXiv:1012.3982 [hep-th]].
- [2] J. M. Maldacena, “The large N limit of superconformal field theories and supergravity,” *Adv. Theor. Math. Phys.* **2**, 231 (1998), [hep-th/9711200].
- [3] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, “Gauge theory correlators from non-critical string theory,” *Phys. Lett. B* **428**, 105 (1998), [arXiv:hep-th/9802109].

- [4] E. Witten, “Anti-de Sitter space and holography,” *Adv. Theor. Math. Phys.* **2**, 253 (1998), [arXiv:hep-th/9802150].
- [5] J. A. Minahan and K. Zarembo, “The Bethe ansatz for N=4 super Yang-Mills,” *JHEP* **0303**, 013 (2003) [hep-th/0212208].
- [6] N. Beisert, C. Kristjansen and M. Staudacher, “The Dilatation operator of conformal N=4 super Yang-Mills theory,” *Nucl. Phys. B* **664**, 131 (2003) [hep-th/0303060].
- [7] N. Beisert, “The complete one loop dilatation operator of N=4 super Yang-Mills theory,” *Nucl. Phys. B* **676**, 3 (2004) [hep-th/0307015].
- [8] N. Beisert, “The $su(2|3)$ dynamic spin chain,” *Nucl. Phys. B* **682**, 487 (2004) [hep-th/0310252].
- [9] N. Beisert, “The $SU(2|2)$ dynamic S-matrix,” *Adv. Theor. Math. Phys.* **12**, 945 (2008) [hep-th/0511082].
- [10] I. Bena, J. Polchinski and R. Roiban, “Hidden symmetries of the $AdS(5) \times S^{*5}$ superstring,” *Phys. Rev. D* **69**, 046002 (2004) [hep-th/0305116].
- [11] N. Beisert, B. Eden and M. Staudacher, “Transcendentality and Crossing,” *J. Stat. Mech.* **0701**, P01021 (2007) [hep-th/0610251].
- [12] L. Freyhult, “Review of AdS/CFT Integrability, Chapter III.4: Twist States and the cusp Anomalous Dimension,” *Lett. Math. Phys.* **99**, 255 (2012) [arXiv:1012.3993 [hep-th]].
- [13] O. Aharony, O. Bergman, D. L. Jafferis and J. Maldacena, “N=6 superconformal Chern-Simons-matter theories, M2-branes and their gravity duals,” *JHEP* **0810**, 091 (2008) [arXiv:0806.1218 [hep-th]].
- [14] O. Aharony, O. Bergman and D. L. Jafferis, “Fractional M2-branes,” *JHEP* **0811**, 043 (2008) [arXiv:0807.4924 [hep-th]].
- [15] J. A. Minahan and K. Zarembo, “The Bethe ansatz for superconformal Chern-Simons,” *JHEP* **0809**, 040 (2008), arXiv:0806.3951 [hep-th].
- [16] D. Bak and S. -J. Rey, “Integrable Spin Chain in Superconformal Chern-Simons Theory,” *JHEP* **0810**, 053 (2008) [arXiv:0807.2063 [hep-th]].
- [17] D. Bak, D. Gang and S. -J. Rey, “Integrable Spin Chain of Superconformal $U(M) \times$ anti- $U(N)$ Chern-Simons Theory,” *JHEP* **0810**, 038 (2008) [arXiv:0808.0170 [hep-th]].
- [18] C. Ahn and R. I. Nepomechie, “N=6 super Chern-Simons theory S-matrix and all-loop Bethe ansatz equations,” *JHEP* **0809**, 010 (2008) [arXiv:0807.1924 [hep-th]].
- [19] G. Arutyunov, S. Frolov, “Superstrings on $AdS_4 \times CP^3$ as a Coset Sigma-model,” *JHEP* **0809** (2008) 129, [arXiv:0806.4940 [hep-th]].
- [20] B. Stefanski, jr, “Green-Schwarz action for Type IIA strings on $AdS_4 \times CP^3$,” *Nucl. Phys.* **B808** (2009) 80-87, [arXiv:0806.4948 [hep-th]].
- [21] J. Gomis, D. Sorokin and L. Wulff, “The Complete $AdS_4 \times CP^3$ superspace for the type IIA superstring and D-branes,” *JHEP* **0903**, 015 (2009), [arXiv:0811.1566 [hep-th]].
- [22] D. Sorokin and L. Wulff, “Evidence for the classical integrability of the complete $AdS_4 \times CP^3$ superstring,” *JHEP* **1011**, 143 (2010), [arXiv:1009.3498 [hep-th]].
- [23] D. Sorokin and L. Wulff, “Peculiarities of String Theory on $AdS_4 \times CP^3$,” arXiv:1101.3777 [hep-th].

- [24] D. Gaiotto, S. Giombi and X. Yin, “Spin Chains in N=6 Superconformal Chern-Simons-Matter Theory,” JHEP **0904**, 066 (2009) [arXiv:0806.4589 [hep-th]].
- [25] G. Grignani, T. Harmark and M. Orselli, “The $SU(2) \times SU(2)$ sector in the string dual of $\mathcal{N} = 6$ superconformal Chern-Simons theory,” Nucl. Phys. B **810**, 115 (2009) [arXiv:0806.4959 [hep-th]].
- [26] C. Kristjansen, M. Orselli, K. Zoubos, “Non-planar ABJM Theory and Integrability,” JHEP **0903**, 037 (2009), [arXiv:0811.2150 [hep-th]].
- [27] P. Caputa, C. Kristjansen and K. Zoubos, “Non-planar ABJ Theory and Parity,” Phys. Lett. B **677**, 197 (2009) [arXiv:0903.3354 [hep-th]].
- [28] A. Mauri, A. Santambrogio and S. Scoleri, “The Leading Order Dressing Phase in ABJM Theory,” [arXiv: 1301.7732[hep-th]].
- [29] D. Astolfi, V. G. M. Puletti, G. Grignani, T. Harmark and M. Orselli, “Finite-size corrections in the $SU(2) \times SU(2)$ sector of type IIA string theory on $AdS_4 \times CP^3$,” Nucl. Phys. B **810**, 150 (2009) [arXiv:0807.1527 [hep-th]].
- [30] T. McLoughlin and R. Roiban, “Spinning strings at one-loop in $AdS_4 \times P^3$,” JHEP **0812**, 101 (2008) [arXiv:0807.3965 [hep-th]].
- [31] L. F. Alday, G. Arutyunov and D. Bykov, “Semiclassical Quantization of Spinning Strings in $AdS_4 \times CP^3$,” JHEP **0811**, 089 (2008) [arXiv:0807.4400 [hep-th]].
- [32] C. Krishnan, “ AdS_4/CFT_3 at One Loop,” JHEP **0809**, 092 (2008) [arXiv:0807.4561 [hep-th]].
- [33] T. McLoughlin, R. Roiban and A. A. Tseytlin, “Quantum spinning strings in $AdS_4 \times CP^3$: Testing the Bethe Ansatz proposal,” JHEP **0811**, 069 (2008) [arXiv:0809.4038 [hep-th]].
- [34] M. A. Bandres and A. E. Lipstein, “One-Loop Corrections to Type IIA String Theory in $AdS_4 \times CP^3$,” JHEP **1004**, 059 (2010) [arXiv:0911.4061 [hep-th]].
- [35] D. Bombardelli and D. Fioravanti, “Finite-Size Corrections of the CP^3 Giant Magnons: The Luscher terms,” JHEP **0907**, 034 (2009) [arXiv:0810.0704 [hep-th]].
- [36] D. Astolfi, V. G. M. Puletti, G. Grignani, T. Harmark and M. Orselli, “Full Lagrangian and Hamiltonian for quantum strings on $AdS_4 \times CP^3$ in a near plane wave limit,” JHEP **1004**, 079 (2010) [arXiv:0912.2257 [hep-th]].
- [37] M. C. Abbott, I. Aniceto and D. Bombardelli, “Quantum Strings and the AdS_4/CFT_3 Interpolating Function,” JHEP **1012**, 040 (2010) [arXiv:1006.2174 [hep-th]].
- [38] D. Astolfi, V. G. M. Puletti, G. Grignani, T. Harmark and M. Orselli, “Finite-size corrections for quantum strings on $AdS_4 \times CP^3$,” JHEP **1105**, 128 (2011) [arXiv:1101.0004 [hep-th]].
- [39] M. C. Abbott, I. Aniceto and D. Bombardelli, “Real and Virtual Bound States in Lüscher Corrections for CP^3 Magnons,” J. Phys. A **45**, 335401 (2012) [arXiv:1111.2839 [hep-th]].
- [40] D. Astolfi, G. Grignani, E. Ser-Giacomi and A. V. Zayakin, “Strings in $AdS_4 \times CP^3$: finite size spectrum vs. Bethe Ansatz,” JHEP **1204**, 005 (2012) [arXiv:1111.6628 [hep-th]].
- [41] M. Beccaria, G. Macorini, C. Ratti and S. Valatka, “Semiclassical folded string in $AdS_4 \times CP^3$,” JHEP **1205**, 030 (2012) [Erratum-ibid. **1205**, 137 (2012)] [arXiv:1203.3852 [hep-th]].
- [42] M. Beccaria, G. Macorini, C. Ratti and S. Valatka, “Semiclassical energy of the $AdS_4 \times CP^3$ folded string,” arXiv:1209.3205 [hep-th].

- [43] T. Klose, “Review of AdS/CFT Integrability, Chapter IV.3: $\mathcal{N} = 6$ Chern-Simons and Strings on $AdS_4 \times CP^3$,” *Lett. Math. Phys.* **99**, 401 (2012), [arXiv:1012.3999 [hep-th]].
- [44] A. E. Lipstein, “Integrability of $\mathcal{N} = 6$ Chern-Simons Theory,” [arXiv:1105.3231 [hep-th]].
- [45] T. Nishioka and T. Takayanagi, “On Type IIA Penrose Limit and N=6 Chern-Simons Theories,” *JHEP* **0808**, 001 (2008) [arXiv:0806.3391 [hep-th]].
- [46] D. Berenstein and D. Trancanelli, “S-duality and the giant magnon dispersion relation,” arXiv:0904.0444 [hep-th].
- [47] R. G. Leigh and M. J. Strassler, “Exactly marginal operators and duality in four-dimensional $\mathcal{N} = 1$ supersymmetric gauge theory,” *Nucl. Phys. B* **447**, 95 (1995) [hep-th/9503121].
- [48] R. Roiban, “On spin chains and field theories,” *JHEP* **0409**, 023 (2004) [hep-th/0312218].
- [49] D. Berenstein and S. A. Cherkis, “Deformations of $\mathcal{N} = 4$ SYM and integrable spin chain models,” *Nucl. Phys. B* **702**, 49 (2004) [hep-th/0405215].
- [50] N. Beisert and R. Roiban, “Beauty and the twist: The Bethe ansatz for twisted N=4 SYM,” *JHEP* **0508**, 039 (2005) [hep-th/0505187].
- [51] S. Frolov, “Lax pair for strings in Lunin-Maldacena background,” *JHEP* **0505**, 069 (2005) [hep-th/0503201].
- [52] O. Lunin and J. M. Maldacena, “Deforming field theories with $U(1) \times U(1)$ global symmetry and their gravity duals,” *JHEP* **0505**, 033 (2005) [hep-th/0502086].
- [53] S. A. Frolov, R. Roiban and A. A. Tseytlin, “Gauge-string duality for superconformal deformations of $\mathcal{N} = 4$ super Yang-Mills theory,” *JHEP* **0507**, 045 (2005) [hep-th/0503192].
- [54] H. -Y. Chen and S. P. Kumar, “Precision test of AdS/CFT in Lunin-Maldacena background,” *JHEP* **0603**, 051 (2006) [hep-th/0511164].
- [55] H. -Y. Chen and K. Okamura, “The Anatomy of gauge/string duality in Lunin-Maldacena background,” *JHEP* **0602**, 054 (2006) [hep-th/0601109].
- [56] N. P. Bobev, H. Dimov and R. C. Rashkov, “Semiclassical strings in Lunin-Maldacena background,” *Bulg. J. Phys.* **35**, 274 (2008) [hep-th/0506063].
- [57] S. Ryang, “Rotating strings with two unequal spins in Lunin-Maldacena background,” *JHEP* **0511**, 006 (2005) [hep-th/0509195].
- [58] C. -S. Chu, G. Georgiou and V. V. Khoze, “Magnons, classical strings and beta-deformations,” *JHEP* **0611**, 093 (2006) [hep-th/0606220].
- [59] N. P. Bobev and R. C. Rashkov, “Spiky strings, giant magnons and beta-deformations,” *Phys. Rev. D* **76**, 046008 (2007) [arXiv:0706.0442 [hep-th]].
- [60] D. V. Bykov and S. Frolov, “Giant magnons in TsT-transformed $AdS_5 \times S^5$,” *JHEP* **0807**, 071 (2008) [arXiv:0805.1070 [hep-th]].
- [61] F. Fiamberti, A. Santambrogio, C. Sieg and D. Zanon, “Finite-size effects in the superconformal beta-deformed N=4 SYM,” *JHEP* **0808**, 057 (2008) [arXiv:0806.2103 [hep-th]].
- [62] C. Ahn and P. Bozhilov, “Finite-Size Dyonic Giant Magnons in TsT-transformed $AdS_5 \times S^5$,” *JHEP* **1007**, 048 (2010) [arXiv:1005.2508 [hep-th]].
- [63] N. Gromov and F. Levkovich-Maslyuk, “Y-system and β -deformed N=4 Super-Yang-Mills,” *J. Phys. A* **44**, 015402 (2011) [arXiv:1006.5438 [hep-th]].

- [64] G. Arutyunov, M. de Leeuw and S. J. van Tongeren, “Twisting the Mirror TBA,” JHEP **1102**, 025 (2011) [arXiv:1009.4118 [hep-th]].
- [65] C. Ahn, Z. Bajnok, D. Bombardelli and R. I. Nepomechie, “Twisted Bethe equations from a twisted S-matrix,” JHEP **1102**, 027 (2011) [arXiv:1010.3229 [hep-th]].
- [66] M. de Leeuw and T. Lukowski, “Twist operators in N=4 beta-deformed theory,” JHEP **1104**, 084 (2011) [arXiv:1012.3725 [hep-th]].
- [67] S. Giardino and V. O. Rivelles, “Pulsating Strings in Lunin-Maldacena Backgrounds,” JHEP **1107**, 057 (2011) [arXiv:1105.1353 [hep-th]].
- [68] C. Ahn, Z. Bajnok, D. Bombardelli and R. I. Nepomechie, “TBA, NLO Luscher correction, and double wrapping in twisted AdS/CFT,” JHEP **1112**, 059 (2011) [arXiv:1108.4914 [hep-th]].
- [69] M. de Leeuw and S. J. van Tongeren, “The spectral problem for strings on twisted $AdS_5 \times S^5$,” Nucl. Phys. B **860**, 339 (2012) [arXiv:1201.1451 [hep-th]].
- [70] C. Ahn, D. Bombardelli and M. Kim, “Finite-size effects of β -deformed AdS_5/CFT_4 at strong coupling,” Phys. Lett. B **710**, 467 (2012) [arXiv:1201.2635 [hep-th]].
- [71] K. L. Panigrahi, P. M. Pradhan and P. K. Swain, “Three Spin Spiky Strings in β -deformed Background,” JHEP **1206**, 057 (2012) [arXiv:1203.3057 [hep-th]].
- [72] C. Ahn, M. Kim and B. -H. Lee, “Worldsheet S-matrix of beta-deformed SYM,” arXiv:1211.4506 [hep-th].
- [73] K. Zoubos, “Review of AdS/CFT Integrability, Chapter IV.2: Deformations, Orbifolds and Open Boundaries,” Lett. Math. Phys. **99**, 375 (2012) [arXiv:1012.3998 [hep-th]].
- [74] E. Imeroni, “On deformed gauge theories and their string/M-theory duals,” JHEP **0810**, 026 (2008) [arXiv:0808.1271 [hep-th]].
- [75] M. Schimpf and R. C. Rashkov, “A Note on strings in deformed $AdS_4 \times CP^3$: Giant magnon and single spike solutions,” Mod. Phys. Lett. A **24**, 3227 (2009) [arXiv:0908.2246 [hep-th]].
- [76] C. Ahn and P. Bozhilov, “Finite-size Giant Magnons on $AdS_4 \times CP^3_\gamma$,” Phys. Lett. B **703**, 186 (2011) [arXiv:1106.3686 [hep-th]].
- [77] J. -B. Wu, “Multi-Spin Strings in $AdS_4 \times CP^3$ and its β -deformations,” arXiv:1208.0389 [hep-th].
- [78] C. Ratti, “Notes on Multi-Spin Strings in $AdS_4 \times CP^3$ and its marginal deformations,” arXiv:1211.4694 [hep-th].
- [79] B. I. Zwiebel, “Two-loop Integrability of Planar N=6 Superconformal Chern-Simons Theory,” J. Phys. A **42**, 495402 (2009) [arXiv:0901.0411 [hep-th]].
- [80] J. A. Minahan, W. Schulgin and K. Zarembo, “Two loop integrability for Chern-Simons theories with N=6 supersymmetry,” JHEP **0903**, 057 (2009) [arXiv:0901.1142 [hep-th]].
- [81] N. Gromov and P. Vieira, “The all loop AdS4/CFT3 Bethe ansatz,” JHEP **0901**, 016 (2009) [arXiv:0807.0777 [hep-th]].
- [82] J. A. Minahan, O. Ohlsson Sax and C. Sieg, “Magnon dispersion to four loops in the ABJM and ABJ models,” J. Phys. A **43**, 275402 (2010) [arXiv:0908.2463 [hep-th]].
- [83] J. A. Minahan, O. Ohlsson Sax and C. Sieg, “Anomalous dimensions at four loops in N=6

- superconformal Chern-Simons theories,” Nucl. Phys. B **846**, 542 (2011) [arXiv:0912.3460 [hep-th]].
- [84] M. Leoni, A. Mauri, J. A. Minahan, O. Ohlsson Sax, A. Santambrogio, C. Sieg and G. Tartaglino-Mazzucchelli, “Superspace calculation of the four-loop spectrum in N=6 supersymmetric Chern-Simons theories,” JHEP **1012**, 074 (2010) [arXiv:1010.1756 [hep-th]].
- [85] D. Bak, H. Min and S. -J. Rey, “Integrability of N=6 Chern-Simons Theory at Six Loops and Beyond,” Phys. Rev. D **81**, 126004 (2010) [arXiv:0911.0689 [hep-th]].